arXiv:math-ph/0312057v1 22 Dec 2003

Second order q–difference equations solvable by factorization method

Alina Dobrogowska and Anatol Odzijewicz

Institute of Theoretical Physics, University of Białystok
Lipowa 41, 15-424 Białystok, Poland
E-mail: alaryzko@alpha.uwb.edu.pl and aodzijew@labfiz.uwb.edu.pl

February 7, 2008

Abstract

By solving an infinite nonlinear system of q-difference equations one constructs a chain of q-difference operators. The eigenproblems for the chain are solved and some applications, including the one related to q-Hahn orthogonal polynomials, are discussed. It is shown that in the limit $q \to 1$ the present method corresponds to the one developed by Infeld and Hull.

Contents

1 Introduction 1

2 Factorized chain of the second order q-difference operators 2

3 Eigenvalue problem for the chain of operators 9

4 q–Hahn orthogonal polynomials 14

5 The case of constant weight functions 16

1 Introduction

The discretization of the ordinary differential equations is an important and necessary step toward finding their numerical solutions. In place of the standard discretization based on the arithmetic progression, one can use a not less efficient q-discretization related to geometric progression. This alternative method leads to q-difference equations, which in the limit $q \to 1$ correspond to the original differential equations. The theory of q-difference equations and the related q-special functions theory have a long history (see e.g. [4]). During the last two decades they have been reviewed because of the great success of the theory of quantum groups.

The other crucial way of solving ordinary differential equations is based on the factorization method first used by Darboux [2]. Later the method was rediscovered many times, in particular by the founders of quantum mechanics, see [8] and [11], while studying the Schrödinger equation. We refer to [16] for an exhaustive presentation of the factorization method. In the paper [2], which is now considered to be fundamental, Infeld and Hull summarised the quantum mechanical applications of the method. Fixing an infinite system of Riccati type equations they have constructed a chain of second order differential operators and proposed some method of solving corresponding eigenproblems.

In this paper we construct the chain (2.71) of second order q-difference operators by solving an infinite nonlinear q-difference system. This chain depends on a freely chosen function and a finite number of real parameters. In Section 2 we find a family of eigenvectors for the operators of (2.71). In Section 3 it is shown that q-Hahn orthogonal polynomials, which are q-deformation of the classical orthogonal polynomials, form the family of solutions obtained by our method. Other examples of solutions obtained by the factorization of q-difference equations are presented in Section 4. Finally passing to the limit $q \to 1$ in (3.36), (3.37) we obtain some new families of solutions for second order differential equations.

2 Factorized chain of the second order q-difference operators

In this section we shall consider the sequence of the second order q–difference operators

$$\mathbf{H}_k = Z_k(x)\partial_q Q^{-1}\partial_q + W_k(x)\partial_q + V_k(x)\ , \qquad k \in \mathbb{N}\cup\{0\} \tag{2.1}$$

acting in the Hilbert spaces $\mathcal{H}_k$. By definition $\mathcal{H}_k$ consist of the complex valued functions $\psi : [a,b]_q \to \mathbb{C}$ defined on the q–interval

$$[a,b]_q := \{q^n a : n \in \mathbb{N}\cup\{0\}\} \cup \{q^n b : n \in \mathbb{N}\cup\{0\}\} \tag{2.2}$$

and square–integrable, i.e. $\langle\psi|\psi\rangle_k < +\infty$, with respect to the scalar products

$$\langle\psi|\varphi\rangle_k := \int\limits_{[a,b]_q} \overline{\psi(x)}\varphi(x)\varrho_k(x)d_qx\ . \tag{2.3}$$

Let us recall (see [34]) that by definition the q–derivative is

$$\partial_q\psi(x) = \frac{\psi(x)-\psi(qx)}{(1-q)x}\ , \tag{2.4}$$

and the q–integral on the q–interval $[a,b]_q$ is given by

$$\int\limits_{[a,b]_q} \psi(x)d_qx := \sum_{n=0}^{\infty}(1-q)q^n\left(b\psi(q^nb) - a\psi(q^na)\right)\ . \tag{2.5}$$

If $a = 0$ and $b = \infty$ then

$$\int\limits_0^\infty \psi(x)d_qx := \lim_{n\to\infty}\int\limits_0^{q^{-n}} \psi(x)d_qx = \sum_{n=-\infty}^{\infty}(1-q)q^nf(q^n)\ . \tag{2.6}$$

In the case if $a = -\infty$ and $b = \infty$

$$\int\limits_{-\infty}^\infty \psi(x)d_qx := \lim_{n\to\infty}\int\limits_{-q^{-n}}^{q^{-n}} \psi(x)d_qx = \sum_{n=-\infty}^{\infty}(1-q)q^n\left(f(q^n)+f(-q^n)\right)\ . \tag{2.7}$$

In the limit $q \to 1$ the above definitions correspond to their counterparts in standard calculus.

The scalar products (2.3) are defined by the weight functions $\varrho_k : [a,b]_q \to \mathbb{R}$, which are related by the recursion relations

$$\varrho_{k-1} = \eta_k \varrho_k \tag{2.8}$$

and

$$\varrho_{k-1} = Q\left(B_k \varrho_k\right) \ , \tag{2.9}$$

where η_k, B_k are real valued functions on $[a,b]_q$ and the operator Q is defined by the formula

$$Q\varphi(x) = \varphi(qx) \ . \tag{2.10}$$

For the sake of consistency we need to add the conditions

$$Q\left(B_k \varrho_k\right) = \eta_k \varrho_k \tag{2.11}$$

on the functions η_k and B_k. Additionally we impose the boundary conditions

$$B_k(a)\varrho_k(a) = B_k(b)\varrho_k(b) = 0 \ . \tag{2.12}$$

If we introduce the functions

$$A_k(x) := \frac{B_k(x) - \eta_k(x)}{(1-q)x} \ , \tag{2.13}$$

we can rewrite the formula (2.11) in the form of a q–Pearson equation [13]

$$\partial_q\left(B_k \varrho_k\right) = A_k \varrho_k \ . \tag{2.14}$$

In the limit $q \to 1$, the equation (2.14) corresponds to the Pearson equation which is important for the theory of classical orthogonal polynomials [1].

We say that the operators $\mathbf{H}_k$ admit a factorization if

$$\mathbf{H}_k = \mathbf{A}_k^* \mathbf{A}_k + a_k \tag{2.15}$$

and

$$\mathbf{H}_k = d_{k+1}^{-1}\left(\mathbf{A}_{k+1}\mathbf{A}_{k+1}^* + a_{k+1}\right) \ , \tag{2.16}$$

where the annihilation operators $\mathbf{A}_k : \mathcal{H}_k \to \mathcal{H}_{k-1}$ are of the form

$$\mathbf{A}_k = \partial_q + f_k \tag{2.17}$$

and f_k are real valued functions on the set $[a,b]_q$. The adjoint operators $\mathbf{A}_k^* : \mathcal{H}_{k-1} \to \mathcal{H}_k$, called the creation operators, are given by

$$\mathbf{A}_k^* = \left(\partial_q + f_k\right)^* = B_k\left(-\partial_q Q^{-1} + f_k\right) - A_k\left(1 + (1-q)x f_k\right) \ . \tag{2.18}$$

The derivation of the formula (2.18) is given in Appendix A. It follows from (2.15) that the real valued functions Z_k, W_k and V_k are related to f_k, B_k, A_k by the formulas:

$$Z_k = -B_k Q^{-1}\left(1 + (1-q)id\ f_k\right) \ , \tag{2.19}$$

$$W_k = B_k f_k - A_k\left(1 + (1-q)id\ f_k\right) - q^{-1} B_k Q^{-1}(f_k) \ , \tag{2.20}$$

$$V_k = -B_k \partial_q\left(Q^{-1}(f_k)\right) - A_k f_k\left(1 + (1-q)id\ f_k\right) + B_k f_k^2 + a_k \ . \tag{2.21}$$

Necessary and sufficient conditions for the consistency of factorization formulas (2.15) and (2.16) are

$$\eta_{k+1}(x) = g_k(x)\eta_k(q^{-1}x) \ , \tag{2.22}$$

$$\varphi_{k+1}(x) = \frac{d_{k+1}}{g_k(x)}\varphi_k(q^{-1}x) \ , \tag{2.23}$$

$$\alpha_k(x) - \frac{g_k(qx)}{d_{k+1}}\alpha_k(qx) = \tag{2.24}$$

$$= \left(\frac{q^2 d_{k+1} B_k(qx) - g_k(q^2x) B_k(q^2x)}{(1-q)^2 q^3 x^2} + d_{k+1}a_k - a_{k+1} \right) \frac{g_k(qx)}{d_{k+1}^2} \ ,$$

where we have introduced the additional notations

$$g_k(x) := \frac{B_{k+1}(x)}{B_k(x)} \ , \tag{2.25}$$

$$\varphi_k(x) := f_k(x) + \frac{1}{(1-q)x} \ , \tag{2.26}$$

$$\alpha_k(x) := \varphi_k^2(x)\eta_k(x) \ . \tag{2.27}$$

The detailed derivation of these formulas is given in Appendix B and in the paper [12, 5].

The relations (2.22), (2.23) and (2.25), (2.27) allow us to express the functions B_k, η_k, φ_k and α_k by the initial data B_0, η_0, φ_0 and α_0

$$B_k(x) = g_{k-1}(x) g_{k-2}(x) \dots g_0(x) B_0(x) \ , \tag{2.28}$$

$$\eta_k(x) = g_{k-1}(x) g_{k-2}(q^{-1}x) \dots g_0(q^{-k+1}x)\eta_0(q^{-k}x) \ , \tag{2.29}$$

$$\varphi_k(x) = \frac{d_k \dots d_1}{g_{k-1}(x) \dots g_0(q^{-k+1}x)}\varphi_0(q^{-k}x) \ , \tag{2.30}$$

$$\alpha_k(x) = \frac{(d_k \dots d_1)^2}{g_{k-1}(x) \dots g_0(q^{-k+1}x)}\alpha_0(q^{-k}x) \ . \tag{2.31}$$

Substituting (2.28–2.31) into condition (2.24) we obtain the infinite sequence of the non-linear functional equations

$$\alpha_0(x) - d_{k+1}\frac{G_{k+1}(x)}{G_k(qx)}\alpha_0(qx) = G_{k+1}(x)\left(d_{k+1}a_k - a_{k+1} + \right. \tag{2.32}$$

$$\left. + \frac{q^2 d_{k+1} g_{k-1}(q^{k+1}x) \dots g_0(q^{k+1}x) B_0(q^{k+1}x) - g_k(q^{k+2}x) \dots g_0(q^{k+2}x) B_0(q^{k+2}x)}{(1-q)^2 q^{2k+3} x^2} \right) \ ,$$

where

$$G_k(x) := \frac{g_{k-1}(q^k x) \dots g_0(qx)}{(d_k \dots d_1)^2} \qquad \text{for} \ \ k \in \mathbb{N} \ , \tag{2.33}$$

$$G_0(x) := 1 \ , \tag{2.34}$$

for the functions α_0, B_0 and g_k for $k \in \mathbb{N} \cup \{0\}$.

One sees from (2.28–2.31) that the sequence of functions g_k, $k \in \mathbb{N}$, satisfying (2.32) defines the chain of $q-$ difference operators (2.1) if the first element $\mathbf{H}_0$ of the chain is given. So, the problem of construction of the factorized chain given by (2.15) and (2.16) is equivalent to solving of the system of functional equations (2.32).

Let us now present the limit behaviour of the formulas obtained above when the parameter q tends to 1. It is easy to see that the set $[a,b]_q$ becomes the interval $[a,b]$ in the limit $q \to 1$ and the scalar product turns to be

$$\langle\psi|\varphi\rangle_k = \int_a^b \overline{\psi(x)}\varphi(x)\varrho_k(x)dx \ , \tag{2.35}$$

where the weight function $\varrho_k(x)$ satisfies the Pearson equation

$$\frac{d}{dx}(\varrho_k B_k) = \varrho_k A_k \ , \tag{2.36}$$

with the boundary conditions (2.12). For $q \to 1$ the operator Q goes to the identity operator and $\partial_q \xrightarrow[q\to 1]{} \frac{d}{dx}$. In the limiting case the annihilation and creation operators are of the form

$$\mathbf{A}_k = \frac{d}{dx} + f_k, \tag{2.37}$$

$$\mathbf{A}_k^* = B_k\left(-\frac{d}{dx} + f_k\right) - A_k \tag{2.38}$$

and the operators $\mathbf{H}_k$ are given by

$$\mathbf{H}_k = -B_k\frac{d^2}{dx^2} - A_k\frac{d}{dx} + (f_k^2 - f_k')B_k - f_kA_k + a_k \ . \tag{2.39}$$

The q-difference equation (2.1) tends to the differential equation

$$\left(Z_k(x)\frac{d^2}{dx^2} + W_k(x)\frac{d}{dx} + V_k(x)\right)\psi_k(x) = \lambda_k\psi_k(x) \ , \tag{2.40}$$

where the coefficients are given by

$$Z_k(x) = -B_k(x) \ , \tag{2.41}$$
$$W_k(x) = -A_k(x) \ , \tag{2.42}$$
$$V_k(x) = \left(f_k^2(x) - f_k'(x)\right)B_k(x) - f_k(x)A_k(x) + a_k \ . \tag{2.43}$$

The recurrence transformations (2.22–2.23) for $q \to 1$ tend to

$$B_{k+1} = d_{k+1}B_k \ , \tag{2.44}$$

$$A_{k+1} = d_{k+1}\left(A_k - \frac{d}{dx}B_k\right). \tag{2.45}$$

The sequence of q-difference equations (2.24) tends to the sequence of non–linear differential equations

$$B_k(f_{k+1}^2 - f_k^2 + f_{k+1}' + f_k') - A_k(f_{k+1} - f_k) + 2B_k'f_{k+1} - A_k' + B_k'' = a_k - \frac{a_{k+1}}{d_{k+1}}, \tag{2.46}$$

$k \in \mathbb{N}\cup\{0\}$. The equation (2.46) for $B_k(x) \equiv 1$ and $A_k(x) \equiv 0$ was considered in many papers (see [7, 9, 10, 11, 15, 16]), but nevertheless for these differential–difference equations there is no complete theory. One of the methods for solving of (2.46) is to look for the solutions of (2.46) in the form of infinite series

$$f_k = \sum_{i\in\mathbb{Z}} \tilde{f}_i(x)k^i \tag{2.47}$$

and obtain in this way the conditions on the function $\tilde{f}_i(x)$. The case of solutions given by the finite series were consider by Infeld and Hull [7]. The classification of all factorisable one–dimensional problems is still an open question.

Now, we come back to the general case. Regarding the extreme nonlinearity of the system (2.32), the possibility to solve it is rather out of the question. Therefore, we shall restrict ourselves to the subcase

$$g_k(x) := d_{k+1}q^\gamma \qquad \text{for} \;\; \gamma\in\mathbb{R} \tag{2.48}$$

and consider the system (2.32), which is reduced now to

$$\alpha_0(x) - q^\gamma\alpha_0(qx) = \frac{q^{(k+1)\gamma}}{d_{k+1}\dots d_1}\left(d_{k+1}a_k - a_{k+1}\right) + \tag{2.49}$$

$$+q^{2(k+1)\gamma}Q^{k+1}\frac{q^{2-\gamma}B_0(x)-B_0(qx)}{(1-q)^2qx^2}\ ,$$

as the infinite system of equations on the initial functions B_0 and α_0. Eliminating α_0 from (2.49) we obtain

$$(1-q)^2q^{3-\gamma}d_1^{-1}x^2\left(\frac{q^{k\gamma}}{d_{k+1}\dots d_1}\left(d_{k+1}a_k - a_{k+1}\right) - d_1a_0 + a_1\right) = \tag{2.50}$$

$$= q^{2-\gamma}B_0(qx) - B_0(q^2x) - q^{2k(\gamma-1)}\left(q^{2-\gamma}B_0(q^{k+1}x) - B_0(q^{k+2}x)\right), \qquad k\in\mathbb{N}\ .$$

Now, we shall look for the solution of (2.50) in the form

$$B_0(x) = x^\delta\sum_{n\in\mathbb{Z}} b_nx^n\ , \tag{2.51}$$

where $\delta\in\langle 0,1)$. Substituting (2.51) into (2.50) and comparing the coefficients in front of x^n we obtain the expressions for the $a_k\in\mathbb{R}$

$$a_{k+1} = d_{k+1}\dots d_1q^{-\gamma k}\left(-a_0\frac{[\gamma k]}{[\gamma]} + \frac{a_1}{d_1}\frac{[\gamma(k+1)]}{[\gamma]} - qb_2[\gamma k][\gamma(k+1)\right), \qquad k\in\mathbb{N} \tag{2.52}$$

and the function B_0

$$B_0(x) = b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma}\ , \tag{2.53}$$

where $b_2, b_1, b_0\in\mathbb{R}$. From (2.53) and (2.49) we have:

(i) if $\gamma\neq 0$, then

$$\alpha_0(x) = \frac{q^{\gamma+1}b_2}{(1-q)^2} + \frac{q^\gamma(d_1a_0-a_1)}{(1-q^\gamma)d_1} + hx^{-\gamma} + \frac{q^{1-\gamma}b_0}{(1-q)^2}x^{-2\gamma}\ , \tag{2.54}$$

where $h\in\mathbb{R}$;

(ii) if $\gamma = 0$, then

$$\alpha_0(x) = h \quad \text{and} \quad d_1a_0 = a_1\ , \tag{2.55}$$

where $h\in\mathbb{R}$.

Finally, substituting (2.48) to (2.28–2.31) we find the following transformation formulas

$$B_k(x) = q^{\gamma k} d_k \dots d_1 B_0(x) \ , \tag{2.56}$$
$$\eta_k(x) = q^{\gamma k} d_k \dots d_1 \eta_0(q^{-k}x) \ , \tag{2.57}$$
$$\varphi_k(x) = q^{-\gamma k} \varphi_0(q^{-k}x) \ , \tag{2.58}$$
$$\alpha_k(x) = q^{-\gamma k} d_k \dots d_1 \alpha_0(q^{-k}x) \ , \tag{2.59}$$

where B_0, α_0 are given by (2.53) and (2.54–2.55) respectively. The functions η_0 and $\varphi_0(x)$ are related to A_0 and α_0 by

$$\eta_0(x) = b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma} - (1-q)xA_0(x) \ , \tag{2.60}$$
$$\varphi_0(x) = \sqrt{\frac{\alpha_0(x)}{\eta_0(x)}} \ . \tag{2.61}$$

At the moment, given the functions B_0, α_0, we can use (2.56–2.59), (2.13–2.14), (2.26) and (2.27) in order to express the functions A_k, f_k and ϱ_k:

$$\begin{aligned} A_k(x) =& q^{\gamma k} d_k \dots d_1 \Big(q^{-k} A_0(q^{-k}x) + [-2k] b_2 x + \\ &+ [k(\gamma-2)] b_1 x^{1-\gamma} + [2k(\gamma-1)] b_0 x^{1-2\gamma} \Big) \ , \end{aligned} \tag{2.62}$$
$$f_k(x) = q^{-\gamma k} f_0(q^{-k}x) - \frac{1-q^{k(1-\gamma)}}{(1-q)x} \ , \tag{2.63}$$
$$\varrho_k(x) = \frac{q^{-\frac{\gamma k(k+1)}{2}}}{d_k d_{k-1}^2 \dots d_1^k} \frac{\varrho_0(q^{-k}x)}{\prod_{n=0}^{k-1} \left(b_2 q^{-2n} x^2 + b_1 q^{n(\gamma-2)} x^{2-\gamma} + b_0 q^{2n(\gamma-1)} x^{2-2\gamma} \right)} \tag{2.64}$$

by A_0, f_0 and ϱ_0. From conditions (2.13–2.14), (2.26) and (2.27) we see that the functions A_0, f_0, ϱ_0 are related by

$$\varrho_0(x) = \frac{q^2 b_2 x + b_1 q^{2-\gamma} x^{1-\gamma} + b_0 q^{2(1-\gamma)} x^{1-2\gamma}}{b_2 x + b_1 x^{1-\gamma} + b_0 x^{1-2\gamma} - (1-q) A_0(x)} \varrho_0(qx) \ , \tag{2.65}$$
$$\left(f_0(x) + \frac{1}{(1-q)x} \right)^2 = \frac{\alpha_0(x)}{b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma} - (1-q)xA_0(x)} \ . \tag{2.66}$$

So, further we shall assume that the function $\frac{A_0(0)}{B_0(0)}$ is continuous in 0. Under this assumption we obtain from (2.65) and (2.66)

$$f_0(x) = \sqrt{\frac{\alpha_0(x)}{b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma} - (1-q)xA_0(x)}} - \frac{1}{(1-q)x} \ , \tag{2.67}$$
$$\varrho_0(x) = \frac{1}{b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma}} \prod_{n=0}^{\infty} Q^n \left(\frac{1}{1 - (1-q)x \frac{A_0(x)}{b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma}}} \right) \ . \tag{2.68}$$

This means that one finds the explicit formulas for the annihilation and creation operators

$$\mathbf{A}_k = \partial_q - \frac{1}{(1-q)x} + q^{-\gamma k} \sqrt{\frac{\alpha_0(q^{-k}x)}{\eta_0(q^{-k}x)}} \ , \tag{2.69}$$
$$\begin{aligned} \mathbf{A}_k^* =& d_k \dots d_1 \left(-q^{\gamma k} (b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma}) \left(\partial_q Q^{-1} + \frac{1}{(1-q)x} \right) + \right. \\ &\left. + \sqrt{\alpha_0(q^{-k}x) \eta_0(q^{-k}x)} \right) \end{aligned} \tag{2.70}$$

and from this the explicit expression for $\mathbf{H}_k$

$$\begin{aligned}\mathbf{H}_k = d_k \dots d_1 \Bigg(& -(1-q)q^{-1}x^3(b_2 + b_1 x^{-\gamma} + b_0 x^{-2\gamma})\sqrt{\frac{\alpha_0(q^{-(k+1)}x)}{\eta_0(q^{-(k+1)}x)}}\partial_q Q^{-1}\partial_q + \\ & + \left(-q^{-1}x^2(b_2 + b_1 x^{-\gamma} + b_0 x^{-2\gamma})\sqrt{\frac{\alpha_0(q^{-(k+1)}x)}{\eta_0(q^{-(k+1)}x)}} + \sqrt{\alpha_0(q^{-k}x)\eta_0(q^{-k}x)}\right)\partial_q + \\ & + \frac{b_2 + b_1 x^{-\gamma} + b_0 x^{-2\gamma}}{(1-q)^2}\left(q - (1-q)x\sqrt{\frac{\alpha_0(q^{-(k+1)}x)}{\eta_0(q^{-(k+1)}x)}}\right) + \\ & + q^{-\gamma k}\alpha_0(q^{-k}x) - \frac{1}{(1-q)x}\sqrt{\eta_0(q^{-k}x)\alpha_0(q^{-k}x)} + \\ & - q^{-\gamma(k-1)}\left(a_0\frac{[\gamma(k-1)]}{[\gamma]} - \frac{a_1}{d_1}\frac{[\gamma k]}{[\gamma]} + q b_2[\gamma(k-1)][\gamma k]\right)\Bigg) ,\end{aligned} \tag{2.71}$$

which depend only on a function A_0.

The chains of operators $\mathbf{A}_k$, $\mathbf{A}^*_k$ and $\mathbf{H}_k$ appearing in (2.69), (2.70) and (2.71) in the limit $q \to 1$ are given by

$$\mathbf{A}_k = \frac{d}{dx} + f_0(x) + \frac{k(\gamma-1)}{x} , \tag{2.72}$$

$$\mathbf{A}^*_k = d_k \dots d_1 \left(B_0(x)\left(-\frac{d}{dx} + f_0(x) + \frac{k(\gamma-1)}{x}\right) - A_0(x) + k\frac{d}{dx}B_0(x)\right) , \tag{2.73}$$

$$\begin{aligned}\mathbf{H}_k = d_k \dots d_1 \Bigg(& -B_0(x)\frac{d^2}{dx^2} - (A_0(x) - kB_0'(x))\frac{d}{dx} + \\ & + \left(f_0^2(x) - f_0'(x) + \frac{2k(\gamma-1)}{x}f_0(x) + \frac{k(\gamma-1)(k(\gamma-1)+1)}{x^2}\right)B_0(x) - \\ & - \left(f_0(x) + \frac{k(\gamma-1)}{x}\right)(A_0(x) - kB_0'(x)) - a_0(k-1) + \frac{a_1}{d_1}k - b_2\gamma^2 k(k-1)\Bigg) ,\end{aligned} \tag{2.74}$$

where

$$B_k(x) = d_k \dots d_1 B_0(x) , \tag{2.75}$$

$$A_k(x) = d_k \dots d_1 (A_0(x) - k\frac{d}{dx}B_0(x)) , \tag{2.76}$$

$$f_k(x) = f_0(x) + k(\gamma-1)\frac{1}{x} , \tag{2.77}$$

$$\varrho_k(x) = \frac{1}{d_k d_{k-1}^2 \dots d_1^k}\frac{\varrho_0(x)}{B_0^k(x)} \tag{2.78}$$

and the functions B_0, f_0 and ϱ_0 have the form

$$B_0(x) = b_2 x^2 + b_1 x^{2-\gamma} + b_0 x^{2-2\gamma} , \tag{2.79}$$

$$f_0(x) = \begin{cases} \frac{-b_2(\gamma+1)x + \frac{d_1 a_0 - a_1}{\gamma d_1}x - b_1\tilde{h}x^{1-\gamma} - b_0(1-\gamma)x^{1-2\gamma} + A_0(x)}{2(b_2x^2 + b_1x^{2-\gamma} + b_0x^{2-2\gamma})} & \text{dla } \gamma \neq 0 \\ -\frac{\tilde{\alpha}}{2}\frac{1}{x} + \frac{A_0(x)}{2(b_2+b_1+b_0)x^2} & \text{dla } \gamma = 0 \end{cases} , \tag{2.80}$$

$$\varrho_0(x) = \frac{1}{B_0(x)} e^{\int_0^x \frac{A_0(t)}{B_0(t)} dt} \ . \tag{2.81}$$

Summing up we see that the construction presented above gives us the nontrivial chain of Hamiltonians (2.71) parameterised by the freely chosen function A_0 and the real parameters b_0, b_1, b_2, $\tilde{h}$, d_k and γ, $k \in \mathbb{N} \cup \{0\}$.

3 Eigenvalue problem for the chain of operators

We shall be interested in solving the eigenvalue problems

$$\mathbf{H}_k \psi_k = \lambda_k \psi_k \quad \text{for} \quad k \in \mathbb{N} \cup \{0\} \ . \tag{3.1}$$

If the operators $\mathbf{H}_k$ admit the factorization given by (2.15) and (2.16) then the eigenvalue equation (3.1) is equivalent to the two equations

$$\mathbf{A}_k^* \mathbf{A}_k \psi_k = (\lambda_k - a_k) \psi_k \ , \tag{3.2}$$

$$\mathbf{A}_{k+1} \mathbf{A}_{k+1}^* \psi_k = (d_{k+1} \lambda_k - a_{k+1}) \psi_k \ . \tag{3.3}$$

From (3.2) and (3.3) one gets

$$\mathbf{H}_k \mathbf{A}_{k+1}^* \psi_k = d_{k+1} a_k \mathbf{A}_{k+1}^* \psi_k \tag{3.4}$$

if

$$\mathbf{H}_k \psi_k = a_k \psi_k \tag{3.5}$$

or equivalently, if

$$\mathbf{A}_k \psi_k = 0 \ . \tag{3.6}$$

Let us remark here that

$$\mathbf{A}_{k+1} \mathbf{A}_{k+1}^* \psi_k = (d_{k+1} \lambda_k - a_{k+1}) \, \psi_k \tag{3.7}$$

and thus $\mathbf{A}_{k+1}^* \psi_k \in \mathcal{H}_{k+1}$. The formulas (3.7) show also that the application of $\mathbf{A}_{k+1}$ to $\mathbf{A}_{k+1}^* \psi_k$ turns it back to the eigenvector of $\mathbf{H}_k$ proportional to the eigenvector ψ_k. Therefore, in the case when $\lambda_k = a_k$ the eigenvalue problem (3.1) is reduced to the equation (3.6) which is a first rank q–difference equation, i.e.

$$\psi_k(x) = \frac{q^{\gamma k}}{(1-q)x} \sqrt{\frac{\eta_0(q^{-k}x)}{\alpha_0(q^{-k}x)}} \psi_k(qx), \tag{3.8}$$

where B_0 and α_0 are given by (2.53) and (2.54–2.55) respectively. By applying the iteration method to (3.8) we find the solution

$$\psi_k(x) = x^{\xi_k} \prod_{n=0}^{\infty} \frac{q^{\xi_k + \gamma k}}{(1-q) q^n x} \sqrt{\frac{\eta_0(q^{n-k}x)}{\alpha_0(q^{n-k}x)}} \ , \tag{3.9}$$

where admissible choices of the real parameter ξ_k and function A_0 are presented in the table below (Figure 1). $A(x)$ is to be an arbitrary analytic function.

Now, let us answer the question of when the solution ψ_k of (3.9) belongs to the Hilbert space $\mathcal{H}_k$. In order to do this we observe that

$$\left(|\psi_k|^2 \varrho_k \right)(x) = \frac{q^{2\gamma k}}{(1-q)^2 x^2} \frac{B_0(qx)}{\alpha_0(q^{-k}x)} \left(|\psi_k|^2 \varrho_k \right)(qx) \ . \tag{3.10}$$

		$A_0(x)$	ξ_k
$\gamma > 0$	$b_0 \neq 0$	$x^{1-2\gamma}A(x)$	$-(\gamma-1)k - \frac{1}{2}\log_q\left(q^{\gamma-1} - (1-q)q^{\gamma-1}\frac{A(0)}{b_0}\right)$
	$b_0 = 0$ $b_1 \neq 0$ $h \neq 0$	$x^{1-\gamma}A(x)$	$-(\gamma-1)k - \frac{1}{2}\log_q\left(\frac{b_1-(1-q)A(0)}{(1-q)^2 h}\right)$
	$b_0 = b_1 = h = 0$ $b_2 \neq 0$ $b_2 \neq \frac{(1-q)(a_1-d_1a_0)}{[\gamma]qd_1}$	$xA(x)$	$-(\gamma-1)k - \frac{1}{2}\log_q\left(\frac{b_2-(1-q)A(0)}{q^{\gamma+1}b_2+\frac{(1-q)q^{\gamma}(d_1a_0-a_1)}{[\gamma]d_1}}\right)$
$\gamma = 0$		$xA(x)$	$k - \frac{1}{2}\log_q\left(\frac{b_2+b_1+b_0-(1-q)A(0)}{(1-q)^2\alpha}\right)$
$\gamma < 0$	$b_2 \neq 0$	$xA(x)$	-$(\gamma-1)k - \frac{1}{2}\log_q\left(\frac{b_2-(1-q)A(0)}{q^{\gamma+1}b_2+\frac{(1-q)q^{\gamma}(d_1a_0-a_1)}{[\gamma]d_1}}\right)$
	$b_2 = 0$ $b_1 \neq 0$ $h \neq 0$ $d_1a_0 = a_1$	$x^{1-\gamma}A(x)$	$-(\gamma-1)k - \frac{1}{2}\log_q\left(\frac{b_1-(1-q)A(0)}{(1-q)^2 h}\right)$
	$b_2 = b_1 = h = 0$ $b_0 \neq 0$ $d_1a_0 = a_1$	$x^{1-2\gamma}A(x)$	$-(\gamma-1)k - \frac{1}{2}\log_q\left(q^{\gamma-1} - (1-q)q^{\gamma-1}\frac{A(0)}{b_0}\right)$

Figure 1: Table of the forms of the function A_0 and the parameter ξ_k

The equation (3.10) can be written for $\gamma = 0$ in the form

$$\left(|\psi_k|^2\varrho_k\right)(x) = \frac{q^2\left(b_2+b_1+b_0\right)}{(1-q)^2\alpha}\left(|\psi_k|^2\varrho_k\right)(qx)\ , \tag{3.11}$$

and for $\gamma \neq 0$ in the form

$$\left(|\psi_k|^2\varrho_k\right)(x) = \tag{3.12}$$

$$= \frac{q^{1-\gamma}\left(b_2(qx)^{2\gamma}+b_1(qx)^{\gamma}+b_0\right)}{q^{2\gamma}\left(b_2+\frac{(1-q)^2}{(1-q^{\gamma})}\frac{(d_1a_0-a_1)}{qd_1}\right)(q^{-k}x)^{2\gamma}+(1-q)^2q^{\gamma-1}h(q^{-k}x)^{\gamma}+b_0}\left(|\psi_k|^2\varrho_k\right)(qx)\ .$$

We also observe that the function $|\psi_k|^2\varrho_k$ does not depend on $A_0(x)$. Using iteration method, after standard calculations we obtain the classes of solutions of (3.10) described in the following proposition.

Proposition 1 *For the solutions to the equation (3.10), the following cases hold:*
1. For $\gamma = 0$ *we have*

$$\left(|\psi_k|^2\varrho_k\right)(x) = x^r\ , \tag{3.13}$$

where $q^{-r} = \frac{q^2(b_2+b_1+b_0)}{(1-q)^2\alpha}$.
2. For $\gamma \neq 0$ *we have following possibilities:*

(i) *If* $b_0 \neq 0$, $b_2 \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^{\gamma})}\frac{(d_1a_0-a_1)}{qd_1} \neq 0$, *then*

$$\left(|\psi_k|^2\varrho_k\right)(x) = x^{\gamma-1}\frac{\left(\frac{(qx)^{\gamma}}{x_1};q^{\gamma}\right)_\infty\left(\frac{(qx)^{\gamma}}{x_2};q^{\gamma}\right)_\infty}{\left(\frac{(q^{-k}x)^{\gamma}}{y_1};q^{\gamma}\right)_\infty\left(\frac{(q^{-k}x)^{\gamma}}{y_2};q^{\gamma}\right)_\infty}\ . \tag{3.14}$$

(ii) *If* $b_0 \neq 0,\ b_2 \neq 0,\ h \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} = 0,$ *then*

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^{\gamma-1}\frac{\left(\frac{(qx)^\gamma}{x_1};q^\gamma\right)_\infty\left(\frac{(qx)^\gamma}{x_2};q^\gamma\right)_\infty}{\left(\frac{(q^{-k}x)^\gamma}{y_1};q^\gamma\right)_\infty}\ . \tag{3.15}$$

(iii) *If* $b_0 \neq 0,\ b_2 \neq 0,\ h = 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} = 0,$ *then*

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^{\gamma-1}\left(\frac{(qx)^\gamma}{x_1};q^\gamma\right)_\infty\left(\frac{(qx)^\gamma}{x_2};q^\gamma\right)_\infty\ . \tag{3.16}$$

(iv) *If* $b_0 = 0,\ b_1 \neq 0,\ b_2 \neq 0,\ h \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} \neq 0,$ *then*

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^{r}\frac{\left(\frac{(qx)^\gamma}{x_1};q^\gamma\right)_\infty}{\left(\frac{(q^{-k}x)^\gamma}{y_1};q^\gamma\right)_\infty}\ , \tag{3.17}$$

where $q^{-r} = \left|\frac{q^{2+\gamma(k-1)}b_1}{(1-q)^2h}\right|.$

(v) *If* $b_0 = 0,\ b_1 \neq 0,\ b_2 \neq 0,\ h \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} = 0,$ *then*

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^{r}\left(\frac{(qx)^\gamma}{x_1};q^\gamma\right)_\infty\ , \tag{3.18}$$

where $q^{-r} = \left|\frac{q^{2+\gamma(k-1)}b_1}{(1-q)^2h}\right|.$

(vi) *If* $b_0 = h = 0,\ b_1 \neq 0,\ b_2 \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} \neq 0,$ *then*

(a)

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^{r}\frac{\left(\frac{(qx)^\gamma}{x_1};q^\gamma\right)_\infty}{\left(-(q^{-k}x)^\gamma;q^\gamma\right)_\infty\left(-q^\gamma(q^{-k}x)^{-\gamma};q^\gamma\right)_\infty}\ , \tag{3.19}$$

where $q^{-r} = \frac{q^{k\gamma+1}b_1}{q^{2\gamma}\left(b_2+\frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1}\right)} > 0;$

(b)

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^{r}\frac{\left(\frac{(qx)^\gamma}{x_1};q^\gamma\right)_\infty}{\left((q^{-k}x)^\gamma;q^\gamma\right)_\infty\left(q^\gamma(q^{-k}x)^{-\gamma};q^\gamma\right)_\infty}\ , \tag{3.20}$$

where $-q^{-r} = \frac{q^{k\gamma+1}b_1}{q^{2\gamma}\left(b_2+\frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1}\right)} < 0.$

(vii) *If* $b_0 = b_1 = 0,\ b_2 \neq 0,\ h \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} \neq 0,$ *then*

(a)

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^{r}\frac{\left(-x^\gamma;q^\gamma\right)_\infty\left(-q^\gamma x^{-\gamma};q^\gamma\right)_\infty}{\left(\frac{(q^{-k}x)^\gamma}{y_1};q^\gamma\right)_\infty}\ , \tag{3.21}$$

where $q^{-r} = \frac{q^{2+k\gamma}b_2}{(1-q)^2h} > 0;$

(b)

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^r\,\frac{(x^\gamma;q^\gamma)_\infty\,(q^\gamma x^{-\gamma};q^\gamma)_\infty}{\left(\frac{(q^{-k}x)^\gamma}{y_1};q^\gamma\right)_\infty}\ , \tag{3.22}$$

where $-q^{-r} = \frac{q^{2+k\gamma}b_2}{(1-q)^2h} < 0.$

(viii) *If* $b_0 = b_1 = h = 0,\ b_2 \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} \neq 0,$ *then*

$$\left(|\psi_k|^2\,\varrho_k\right)(x) = x^r\ , \tag{3.23}$$

where $q^{-r} = \left|\frac{q^{1-\gamma+2k\gamma}b_2}{b_2+\frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1}}\right|.$

In all the above cases x_1, x_2 *are roots of the polynomial*

$$b_2x^2 + b_1x + b_0 = 0 \tag{3.24}$$

and y_1, y_2 *are roots of the polynomial*

$$\left(q^{2\gamma}b_2 + (1-q)^2\frac{q^{2\gamma-1}(d_1a_0-a_1)}{(1-q^\gamma)d_1}\right)x^2 + (1-q)^2q^{\gamma-1}hx + b_0 = 0\ . \tag{3.25}$$

Proof: We easily obtain the subcases **(i)** − **(iii)** by iteration. The other cases are proved by calculation of the Laurent expression coefficient and application of Jacobi's identities

$$\sum_{k=-\infty}^{\infty} q^{k^2}x^k = \left(q^2;q^2\right)_\infty\left(-qx;q^2\right)_\infty\left(-q/z;q^2\right)_\infty\ , \tag{3.26}$$

(see [4]).

□

The proposition given below classifies those function (3.9) which are elements of Hilbert space $\mathcal{H}_k$.

Proposition 2 *The solution (3.9) of equation (3.6) belongs to the Hilbert space* $\mathcal{H}_k$ *if and only if the parameters* $b_0,\ b_1,\ b_2,\ \alpha,\ h,\ d_1,\ a_0,\ a_1$ *and* γ *satisfy the following conditions:*

1. $\gamma = 0$ *and* $\frac{\alpha}{b_2+b_1+b_0} < \frac{q}{(1-q)^2}$.

2. $\gamma > 0$ *and one the following conditions is fulfilled:*
   - **(i)** $b_0 \neq 0,\ b_2 \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} \neq 0$*;*
   - **(ii)** $b_0 \neq 0,\ b_2 \neq 0,\ h \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} = 0$*;*
   - **(iii)** $b_0 \neq 0,\ b_2 \neq 0,\ h = 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} = 0$*;*
   - **(iv)** $b_0 = 0,\ b_1 \neq 0,\ b_2 \neq 0,\ h \neq 0,\ b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} \neq 0$ *and* $\frac{h}{b_1} < \frac{q^{1+\gamma(k-1)}}{(1-q)^2}$*;*
   - **(v)** $b_0 = 0,\ b_1 \neq 0,\ b_2 \neq 0,\ h \neq 0,\ b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} = 0$ *and* $\frac{h}{b_1} < \frac{q^{1+\gamma(k-1)}}{(1-q)^2}$*;*
   - **(vi)** $b_0 = h = 0,\ b_1 \neq 0,\ b_2 \neq 0$ *and* $b_2 + \frac{(1-q)^2}{(1-q^\gamma)}\frac{(d_1a_0-a_1)}{qd_1} \neq 0$*;*
   - **(vii)** *in this case the solutions never belong to the Hilbert space;*

(viii) $b_0 = b_1 = h = 0$, $b_2 \neq 0$, $b_2 + \frac{(1-q)^2}{(1-q^\gamma)} \frac{(d_1 a_0 - a_1)}{q d_1} \neq 0$ *and* $\frac{d_1 a_0 - a_1}{q d_1 b_2} < \frac{1-q^\gamma}{(1-q)^2} \left(q^{\gamma(2k-1)} - 1\right)$.

The notation and classification given above are compatible with Proposition 1.

Proof: The function ψ_k belongs to the Hilbert space if

$$\int\limits_{[a,b]_q} \left(|\psi_k|^2 \varrho_k \right)(x) d_q x < +\infty \,. \tag{3.27}$$

This is equivalent to

$$\sum_{n=0}^{\infty} (1-q) q^n y \left(\left|\psi_k^0\right|^2 \varrho_k \right)(q^n y) < +\infty \tag{3.28}$$

for $y = a, b$. So, for the case **(i)** (i.e. $b_0 \neq 0$, $b_2 \neq 0$ and $b_2 + \frac{(1-q)^2}{(1-q^\gamma)} \frac{(d_1 a_0 - a_1)}{q d_1} \neq 0$) we have from Proposition 1 that the $\left|\psi_k^0\right|^2 \varrho_k$ is given by (3.14), and we show that

$$(1-q) y^\gamma \sum_{n=0}^{\infty} q^{\gamma n} \frac{\left(\frac{(q^{n+1} y)^\gamma}{x_1}; q^\gamma \right)_\infty \left(\frac{(q^{n+1} y)^\gamma}{x_2}; q^\gamma \right)_\infty}{\left(\frac{(q^{n-k} y)^\gamma}{y_1}; q^\gamma \right)_\infty \left(\frac{(q^{n-k} y)^\gamma}{y_2}; q^\gamma \right)_\infty} < +\infty \,. \tag{3.29}$$

From the identity

$$(q^{n\gamma} a; q^\gamma)_\infty = \frac{(a; q^\gamma)_\infty}{(a; q^\gamma)_n} \,, \tag{3.30}$$

where

$$(a; q^\gamma)_\infty = (1-a)(1-q^\gamma a) \ldots \,, \tag{3.31}$$

$$(a; q^\gamma)_n = (1-a)(1-q^\gamma a) \ldots (1 - q^{\gamma(n-1)} a) \,, \tag{3.32}$$

we obtain the conditions equivalent to (3.29)

$$(1-q) y^\gamma \frac{\left(\frac{(qy)^\gamma}{x_1}; q^\gamma \right)_\infty \left(\frac{(qy)^\gamma}{x_2}; q^\gamma \right)_\infty}{\left(\frac{(q^{-k} y)^\gamma}{y_1}; q^\gamma \right)_\infty \left(\frac{(q^{-k} y)^\gamma}{y_2}; q^\gamma \right)_\infty} \times \tag{3.33}$$

$$\times \sum_{n=0}^{\infty} q^{\gamma n} \frac{\left(\frac{(q^{-k} y)^\gamma}{y_1}; q^\gamma \right)_n \left(\frac{(q^{-k} y)^\gamma}{y_2}; q^\gamma \right)_n}{\left(\frac{(qy)^\gamma}{x_1}; q^\gamma \right)_n \left(\frac{(qy)^\gamma}{x_2}; q^\gamma \right)_n} < +\infty \,.$$

Those conditions are fulfilled for $\gamma > 0$. The proofs of the other cases are similar to the one above.

$\square$

Finally let us come back to the general situation and observe that (3.4), (3.5) and (3.6) imply that the function

$$\psi_k^n(x) := \mathbf{A}_k^* \ldots \mathbf{A}_{k-n+1}^* \psi_{k-n}^0(x) \,, \qquad n = 1, \ldots, k, \tag{3.34}$$

is an eigenvector of the operator $\mathbf{H}_k$ with the eigenvalue

$$\lambda_k^n = d_k d_{k-1} \ldots d_{k-n+1} a_{k-n} \tag{3.35}$$

if $\psi_{k-n}^0 := \psi_{k-n}$ is the eigenvector of $\mathcal{H}_{k-n}$ with eigenvalue a_{k-n}. Moreover, one comes back to the eigensubspace $\mathbb{C}\psi_{k-n}^0$ acting on $\mathbb{C}\psi_k^n$ by the annihilation operators $\mathbf{A}_{k-n+1}$, … and



Figure 2: Presentation of action of the operators $\mathbf{A}_k^*$

$\mathbf{A}_k$. The above described procedures can be illustrated by a lattice of points in the (k, n) plane (Figure 2).

The eigenfunctions of the operator $\mathbf{H}_k$ given by (3.9) and (3.34) in the limit $q \to 1$ tend to

$$\psi_k^0(x) = x^{-k(\gamma-1)} e^{-\int_0^x f_0(t)dt} \ , \tag{3.36}$$

$$\psi_k^n(x) = \mathbf{A}_k^* \dots \mathbf{A}_{k-n+1}^* x^{-(k-n)(\gamma-1)} e^{-\int_0^x f_0(t)dt} \quad \text{for} \quad n = 1, 2, \dots k \ , \tag{3.37}$$

with the eigenvalues

$$\lambda_k^n = d_k \dots d_1 \left(-a_0(k-n-1) + \frac{a_1}{d_1}(k-n) - b_2\gamma^2(k-n)(k-n-1) \right) \ . \tag{3.38}$$

In the next sections we want to present some important examples, including the example of orthogonal polynomials of q–Hahn class which, in the limit $q \to 1$, gives classical orthogonal polynomials. These examples will illustrate how the factorization method presented above works.

4 q–Hahn orthogonal polynomials

We obtain q–Hahn orthogonal polynomials when we require that the functions $f_k(x) \equiv 0$ and $d_k = q^{-1}$ for $k \in \mathbb{N} \cup \{0\}$. This is equivalent to

$$\gamma = 1 \ , \tag{4.1}$$

$$B_k(x) = B_0(x) = b_2x^2 + b_1x + b_0 \ , \tag{4.2}$$

$$A_0(x) = \left([2]b_2 - qa_0 + q^2a_1\right) x + \frac{b_1}{1-q} - (1-q)h \ . \tag{4.3}$$

We see that the functions B_k and A_0 are second and a first order polynomials respectively. From (2.62) we obtain that the function A_k is also first order polynomial

$$A_k(x) = q^{-k}A_0(q^{-k}x) + \frac{1-Q^{-k}}{(1-q)x}B_0(x) = \widetilde{a_k}x + \widetilde{b_k} \ , \tag{4.4}$$

where

$$\widetilde{a_k} = -q^{-2(k-1)}\left([2(k-1)]b_2 + q^{-1}a_0 - a_1\right)\ , \tag{4.5}$$

$$\widetilde{b_k} = \frac{b_1}{1-q} - (1-q)q^{-k}h\ . \tag{4.6}$$

Hence, the annihilation and creation operators are given by

$$\mathbf{A}_k = \partial_q\ , \tag{4.7}$$

$$\mathbf{A}^*_k = -(b_2x^2 + b_1x + b_0)\partial_q Q^{-1} - \widetilde{a_k}x - \widetilde{b_k} \tag{4.8}$$

and the Hamiltonian by

$$\begin{aligned} H_k = &-(b_2x^2 + b_1x + b_0)\partial_q Q^{-1}\partial_q - \left(\widetilde{a_k}x + \widetilde{b_k}\right)\partial_q + \\ &+ q^{-2(k-1)}\left(-q^{-1}a_0[k-1] + a_1[k] - b_2[k-1][k]\right)\ . \end{aligned} \tag{4.9}$$

The eigenvalue problem for the Hamiltonian (4.9) is known as the q–Hahn equation [6, 13]

$$\left(B_0(x)\partial_q Q^{-1}\partial_q + A_k(x)\partial_q\right)\psi^n_k = \lambda^n_k\psi^n_k\ . \tag{4.10}$$

The eigenvectors related to the eigenvalues

$$\lambda^0_k = 0\ , \tag{4.11}$$

$$\lambda^n_k = \widetilde{a_k}[n] + b_2[n][n-1]q^{-(n-1)}\ . \tag{4.12}$$

are given by

$$\psi^0_k = 1\ , \tag{4.13}$$

$$\psi^n_k = \mathbf{A}^*_k\dots\mathbf{A}^*_{k-n+1}1 = \prod_{i=k-n+1}^{k}\left(-(b_2x^2 + b_1x + b_0)\partial_q Q^{-1} - \widetilde{a_i}x - \widetilde{b_i}\right)1\ , \tag{4.14}$$

for $k \in \mathbb{N}\cup\{0\}$ and $n = 1, 2, \dots, k$. The functions ψ^0_k (4.14) are polynomials. Each of the families $\{\psi^n_k\}^k_{n=0}$ is a system of polynomials orthogonal with respect to the scalar product given by Jackson's integral

$$\int\limits_{[a,b]_q} \psi^n_k(x)\psi^m_k(x)\varrho_k(x)d_qx \sim \delta_{nm}\ , \tag{4.15}$$

where the weight functions are obtained from (2.68)

$$\varrho_k(x) = \frac{\varrho_0(q^{-k}x)}{B_0(q^{-k+1}x)\dots B_0(x)}\ . \tag{4.16}$$

The classes of the weight functions ϱ_0 and the set of integration $[a,b]_q$ in (4.15) are presented in [13].

In the limit $q \to 1$ this case gives us the classical orthogonal polynomials

$$\left(B_0(x)\frac{d^2}{dx^2} + A_k(x)\frac{d}{dx}\right)P^n_k(x) = \lambda^n_k P^n_k(x)\ . \tag{4.17}$$

The functions B_0 and A_k are second and first order polynomials given by

$$B_k(x) = B_0(x) = b_2x^2 + b_1x + b_0\ , \tag{4.18}$$

$$A_k(x) = \tilde{a}_k x + \tilde{b}_k \; , \tag{4.19}$$

where

$$\tilde{a}_k = -2(k-1)b_2 + a_1 - a_0 \; , \tag{4.20}$$

$$\tilde{b}_k = b_1(\tilde{h} - k) \; , \tag{4.21}$$

(in order to obtain this formulas we demand additionally that $h = \frac{b_1 q^{\tilde{h}}}{(1-q)^2}$ in (4.6)). The eigenvectors ψ_k^n (orthogonal polynomials), in the limiting case, have the forms

$$\psi_k^0(x) = 1 \; , \tag{4.22}$$

$$\psi_k^n(x) = \left(B_0(x)\frac{d}{dx} + A_k(x)\right)\left(B_0(x)\frac{d}{dx} + A_{k-1}(x)\right)\ldots\left(B_0(x)\frac{d}{dx} + A_{k-n+1}(x)\right)1 \tag{4.23}$$

and correspond to the eigenvalues

$$\lambda_k^n = \tilde{a}_k n + b_2 n(n-1) \; . \tag{4.24}$$

5 The case of constant weight functions

We assume that all weight functions are constant $\varrho_k(x) \equiv \text{const}$. We obtain two cases, which we consider below

q–Deformation of the harmonic oscillator
Additionally we demand that $d_k = q^{-1}$ i $b_0 = \varrho_0 = 1$ for the sake of transparency of the formulas. In this case we have:

$$\begin{aligned}
&\gamma = 1 \; , && (5.1)\\
&B_k(x) = 1 \; , && (5.2)\\
&A_k(x) = 0 \; , && (5.3)\\
&f_k(x) = q^{-k} f_0(q^{-k}x) \; , && (5.4)\\
&\varrho_k = 1 \; , && (5.5)
\end{aligned}$$

where

$$f_0(x) = \sqrt{\frac{q^2(q^{-1}a_0 - a_1)}{1-q} + \frac{h}{x} + \frac{1}{(1-q)^2}\frac{1}{x^2}} - \frac{1}{(1-q)x} \; . \tag{5.6}$$

The annihilation and creation operators are given by

$$\begin{aligned}
\mathbf{A}_k &= \partial_q + q^{-k} f_0(q^{-k}x) \; , && (5.7)\\
\mathbf{A}_k^* &= -\partial_q Q^{-1} + q^{-k} f_0(q^{-k}x) \; . && (5.8)
\end{aligned}$$

Solving the equation (3.8) we find the basic state ψ_k^0 of the Hamiltonian given by

$$\begin{gathered}
\mathbf{H}_k = -\left(1 + (1-q)q^{-k-1} x f_0(q^{-k-1}x)\right)\partial_q Q^{-1}\partial_q + \qquad (5.9)\\
+q^{-k}\left(f_0(q^{-k}x) - q^{-1} f_0(q^{-k-1}x)\right)\partial_q +\\
-q^{-k}\partial_q(f_0(q^{-k-1}x)) + q^{-2k} f_0^2(q^{-k}x) + q^{-2k}\left(a_0 + (q^2 a_1 - a_0)[k]\right) \; .
\end{gathered}$$

1. If $a_0 \neq qa_1$, then
$$\psi_k^0(x) = \frac{C_k^0}{\sqrt{\left(\frac{q^{-k}x}{x_1};q\right)_\infty \left(\frac{q^{-k}x}{x_2};q\right)_\infty}} \ , \tag{5.10}$$
where x_1 and x_2 are roots of the polynomial
$$(1-q)q^2(q^{-1}a_0 - a_1)x^2 + (1-q)^2hx + 1 = 0 \tag{5.11}$$
and $C_k^0 \in \mathbb{R} \setminus \{0\}$.

2. If $a_0 = qa_1$ i $h \neq 0$, then
$$\psi_k^0(x) = \frac{C_k^0}{\sqrt{(-(1-q)^2hq^{-k}x;q)_\infty}} \ . \tag{5.12}$$

It easy to see that the operator Q^{-1} acts as follows

$$\begin{tikzcd} \psi_0^0 \arrow[r, "\frac{C_1^0}{C_0^0}Q^{-1}"] & \psi_1^0 \arrow[r, "\frac{C_2^0}{C_1^0}Q^{-1}"] & \cdots \arrow[r, "\frac{C_k^0}{C_{k-1}^0}Q^{-1}"] & \psi_k^0 \arrow[r, "\frac{C_{k+1}^0}{C_k^0}Q^{-1}"] & \cdots \ , \end{tikzcd}$$

or equivalently
$$\psi_k^0(x) = Q^{-k}\psi_0^0(x) \ . \tag{5.13}$$

The functions ψ_k^0 are eigenvectors of the Hamiltonians $\mathbf{H}_k$ with the eigenvalues
$$\lambda_k^0 = a_k = q^{-2k}\left(a_0 + (q^2a_1 - a_0)[k]\right) \ . \tag{5.14}$$

Similarly it is easy to obtain that the functions
$$\psi_k^n(x) = Q^{-k}\psi_0^n(x) \tag{5.15}$$
are eigenvectors of $\mathbf{H}_k$ with
$$\lambda_k^n = q^{-2k}\left(\lambda_0^n + (q^2a_1 - a_0)[k]\right) \ , \tag{5.16}$$
in view the following commutation relations
$$q\mathbf{A}_k^*Q^{-1} = Q^{-1}\mathbf{A}_{k-1}^* \ , \tag{5.17}$$
$$\mathbf{A}_k^*Q = qQ\mathbf{A}_{k+1}^* \ , \tag{5.18}$$
$$q\mathbf{A}_kQ^{-1} = Q^{-1}\mathbf{A}_{k-1} \ , \tag{5.19}$$
$$\mathbf{A}_kQ = qQ\mathbf{A}_{k+1} \ . \tag{5.20}$$

Finally we present the action of the operators diagrammatically Figure 3 and state the following

Proposition 3 *The functions*
$$\psi_k^n(x) = \frac{1}{\sqrt{(a_0 - qa_1)^n n_q! q^{n(n-1)+k}}} Q^{n-k}\mathbf{A}_n^* \dots \mathbf{A}_1^*\psi_0^0(x) \ , \tag{5.21}$$
for $k \in \mathbb{N} \cup \{0\}$ and $n \in \mathbb{N} \cup \{0\}$, where the function ψ_0^0 is given by (5.10) or (5.12), are the eigenvectors of Hamiltonians (5.9) corresponding to the eigenvalues
$$\lambda_k^n = q^{-2k+n}\left(a_0 + (q^2a_1 - a_0)[k-n]\right) \ . \tag{5.22}$$



Figure 3: Presentation of action of the operators

In the limit $q \to 1$ this case gives us the harmonic oscillator

$$\mathbf{H}_k = -\frac{d^2}{dx^2} + \frac{(a_0 - a_1)^2}{4} x^2 + \frac{a_1 + a_0}{2} + (a_1 - a_0)k \ . \tag{5.23}$$

with eigenvectors

$$\psi_k^n(x) = \left(-\frac{d}{dx} + \frac{a_0 - a_1}{2} x \right)^n e^{-\frac{a_0 - a_1}{4} x^2} \quad \text{for} \quad n \in \mathbb{N} \cup \{0\} \tag{5.24}$$

corresponding to the eigenvalues

$$\lambda_k^n = a_0 + (a_0 - a_1)(n - k) \ . \tag{5.25}$$

q–**Deformation of the three–dimensional isotropic harmonic oscillator**
Additionally we demand that $d_k = q^{-2}$ and $b_1 = \varrho_0 = 1$. In this case we have

$$\gamma = 2 \ , \tag{5.26}$$

$$B_k(x) = 1 \ , \tag{5.27}$$

$$A_k(x) = 0 \ , \tag{5.28}$$

$$f_k(x) = q^{-2k} f_0(q^{-k} x) - \frac{1 - q^{-k}}{(1 - q)x} \ , \tag{5.29}$$

$$\varrho_k = 1 \ , \tag{5.30}$$

where

$$f_0(x) = \sqrt{\frac{q^4(q^{-2} a_0 - a_1)}{1 - q^2} + \frac{h}{x^2}} - \frac{1}{(1 - q)x} \ . \tag{5.31}$$

The annihilation and creation operators have the form

$$\mathbf{A}_k = \partial_q + q^{-2k} f_0(q^{-k}x) - \frac{1-q^{-k}}{(1-q)x} \ , \tag{5.32}$$

$$\mathbf{A}^*_k = -\partial_q Q^{-1} + q^{-2k} f_0(q^{-k}x) - \frac{1-q^{-k}}{(1-q)x} \ , \tag{5.33}$$

and the Hamiltonians are given by the formulas

$$\mathbf{H}_k = -\left(q^{-k} + (1-q)q^{-2k-1}xf_0(q^{-k-1}x)\right)\partial_q Q^{-1}\partial_q + \tag{5.34}$$

$$+q^{-2k}\left(f_0(q^{-k}x) - q^{-1}f_0(q^{-k-1}x)\right)\partial_q - q^{-2k}\left(\partial_q f_0(q^{-k-1}x)\right) + \frac{q^{-2k}[k][k+1]}{x^2} +$$

$$+q^{-4k}f_0^2(q^{-k}x) + 2q^{-3k}\frac{[k]}{x}f_0(q^{-k}x) + q^{-4k}\left(a_0 + (q^4a_1 - a_0)\frac{[2k]}{[2]}\right) \ ,$$

The basic states of the Hamiltonians (5.34) can be found as the solution (3.8).

1. If $a_0 \neq q^2 a_1$, then

$$\psi^0_k(x) = \frac{C^0_k}{\sqrt{\left(-\frac{q^4(q^{-2}a_0-a_1)}{(1-q^2)h}q^{-2k}x^2;q\right)_\infty}}x^{\xi_k} \ , \tag{5.35}$$

   where $C^0_k \in \mathbb{R}\setminus\{0\}$ and

$$\xi_k = -k + \log_q(1-q)\sqrt{h} \ . \tag{5.36}$$

2. If $a_0 = q^2 a_1$, then

$$\psi^n_k(x) = C^0_k x^{\xi_k} \ . \tag{5.37}$$

These are the eigenfunctions of the Hamiltonian corresponding to the eigenvalues

$$\lambda^0_k = a_k = q^{-4k}\left(a_0 + (q^4a_1 - a_0)\frac{[2k]}{[2]}\right) \ . \tag{5.38}$$

Finally we have the following lemma:

Proposition 4 *The functions*

$$\psi^n_k(x) = \mathbf{A}^*_k \ldots \mathbf{A}^*_{k-n+1}\psi^0_{k-n} = \tag{5.39}$$

$$= \prod_{i=k-n+1}^{k}\left(\frac{1}{(1-q)x}\left(-Q^{-1} + q^{-k}(1-q)\sqrt{h}\sqrt{1+\frac{q^4(q^{-2}a_0-a_1)}{(1-q^2)h}q^{-2k}x^2}\right)\right)\psi^0_{k-n} \ ,$$

for $n = 1, 2, \ldots, k$, are the eigenvectors of the Hamiltonian with the eigenvalues

$$\lambda^n_k = q^{-2n}a_{k-n} = q^{-2(2k-n)}\left(a_0 + (q^4a_1 - a_0)\frac{[2(k-n)]}{[2]}\right) \ . \tag{5.40}$$

In the limit $q \to 1$ this case gives us the three–dimensional isotropic harmonic oscillator

$$\mathbf{H}_k = -\frac{d^2}{dx^2} + \frac{(k-\frac{\tilde{h}}{2})(k-\frac{\tilde{h}}{2}+1)}{x^2} + \frac{(a_0-a_1)^2}{16}x^2 - \tag{5.41}$$

$$-\frac{a_0-a_1}{2}(k+\frac{\tilde{h}}{2})+\frac{3a_0+a_1}{4}\ .$$

with eigenvectors

$$\psi_k^0(x)=C_k^0 x^{\frac{\tilde{h}}{2}-k}e^{-\frac{a_0-a_1}{8}x^2}\ , \tag{5.42}$$

$$\psi_k^n(x)=\prod_{i=k-n+1}^{k}\left(-\frac{d}{dx}+\frac{a_0-a_1}{4}x-\frac{\tilde{h}}{2}\frac{1}{x}+\frac{i}{x}\right)x^{\frac{\tilde{h}}{2}-k}e^{-\frac{a_0-a_1}{8}x^2}\quad \text{for}\quad n=1,\dots,k \tag{5.43}$$

corresponding to the eigenvalues

$$\lambda_k^n=a_0+(a_1-a_0)(k-n)\ . \tag{5.44}$$

Appendix A. Derivation of formula (2.18)

By the definition of the adjoint operator we have

$$\langle Q^*\psi_k|\varphi_k\rangle_k=\langle\psi_k|Q\varphi_k\rangle_k=\int_a^b\overline{\psi_k(x)}\varphi_k(qx)\varrho_k(x)d_qx=$$

$$=\sum_{n=0}^{\infty}(1-q)q^n b\overline{\psi_k(q^nb)}\varphi_k(q^{n+1}b)\varrho_k(q^nb)-\sum_{n=0}^{\infty}(1-q)q^n a\overline{\psi_k(q^na)}\varphi_k(q^{n+1}a)\varrho_k(q^na)=$$

$$\overset{m=n+1}{=}\sum_{m=1}^{\infty}(1-q)q^m bq^{-1}\overline{\psi_k(q^{m-1}b)}\varphi_k(q^mb)\varrho_k(q^{m-1}b)-$$

$$-\sum_{m=1}^{\infty}(1-q)q^m aq^{-1}\overline{\psi_k(q^{m-1}a)}\varphi_k(q^ma)\varrho_k(q^{m-1}a)\ .$$

In this sum the expression for $m=0$, i.e.

$$(1-q)\left(b\overline{\psi_k(q^{-1}b)}\varphi_k(b)\varrho_k(b)-a\overline{\psi_k(q^{-1}a)}\varphi_k(a)\varrho_k(a)\right)\ , \tag{5.45}$$

does not appear. The functions $\psi_k(x)$ i $\varphi_k(x)$ are defined on the set $\{q^nb:n\in\mathbb{N}\cup\{0\}\}\cup\{q^na:n\in\mathbb{N}\cup\{0\}\}$ and for the other points we shall put these functions equal to zero

$$\begin{array}{c}\left(Q^{-1}\psi\right)(b)=0\ ,\\ \left(Q^{-1}\psi\right)(a)=0\ .\end{array}$$

From the equation (2.11) we obtain for $x\neq a$ and $x\neq b$ that

$$\langle Q^*\psi_k|\varphi_k\rangle_k=\int_a^b\overline{\psi_k(q^{-1}x)}\varphi_k(x)\varrho_k(x)\frac{B_k(x)}{\eta_k(q^{-1}x)}q^{-1}d_qx=\left\langle q^{-1}\frac{B_k}{Q^{-1}\eta_k}(Q^{-1}\psi_k)\middle|\varphi_k\right\rangle_k\ . \tag{5.46}$$

Similarly we have

$$\langle f\psi_k|\varphi_{k-1}\rangle_{k-1}=\int_a^b\overline{f(x)\psi_k(x)}\varphi_{k-1}(x)\varrho_{k-1}(x)d_qx=$$

$$= \int\limits_a^b \overline{\psi_k(x) f(x)} \varphi_{k-1}(x) \eta_k(x) \varrho_k(x) d_q x = \langle \psi_k | \overline{f} \eta_k \varphi_{k-1} \rangle_k \ , \tag{5.47}$$

where we use the equation (2.8).

Summarising we obtain the formula (2.18)

$$\mathbf{A}_k^* = (\partial_q + f_k)^* = B_k \left(-\partial_q Q^{-1} + f_k \right) - A_k \left(1 + (1-q) x f_k \right) \ , \tag{5.48}$$

where the operator Q^{-1} is given by

$$Q^{-1}\varphi(x) = \begin{cases} \varphi(q^{-1}x) & \text{dla} \quad x \neq a \text{ i } x \neq b \\ 0 & \text{dla} \quad x = a \text{ lub } x = b \end{cases} \ . \tag{5.49}$$

Appendix B. Derivation of formulas (2.22–2.24)

The operators of annihilation and creation given by (2.17, 2.18) can be rewritten in the form

$$\mathbf{A}_k = \partial_q + f_k = -\frac{1}{(1-q)x} Q + \varphi_k \ , \tag{5.50}$$

$$\mathbf{A}_k^* = B_k \left(-\partial_q Q^{-1} + f_k \right) - A_k \left(1 + (1-q) x f_k \right) = -\frac{B_k}{(1-q)x} Q^{-1} + \eta_k \varphi_k \ , \tag{5.51}$$

where the functions φ_k, η_k are defined by (2.26) and (2.13). From the conditions (2.15–2.16) we have that

$$\begin{cases} \eta_k(qx)\varphi_k(qx) = d_k \eta_{k-1}(x) \varphi_{k-1}(x) \\ B_k(x)\varphi_k(x) = d_k B_{k-1}(x) \varphi_{k-1}(q^{-1}x) \\ \eta_k(x)\varphi_k^2(x) - d_k \eta_{k-1}(x)\varphi_{k-1}^2(x) = d_k a_{k-1} - a_k + \frac{q^2 d_k B_{k-1}(x) - B_k(qx)}{(1-q)^2 q x^2} \end{cases} . \tag{5.52}$$

The first and second equations of (5.52) are equivalent to

$$\frac{\varphi_k(qx)}{\varphi_{k-1}(x)} = d_k \frac{\eta_{k-1}(x)}{\eta_k(qx)} \ , \tag{5.53}$$

$$\frac{\varphi_k(qx)}{\varphi_{k-1}(x)} = d_k \frac{B_{k-1}(qx)}{B_k(qx)} \ . \tag{5.54}$$

A simple calculation gives us

$$\eta_k(x) = \frac{B_k(x)}{B_{k-1}(x)} \eta_{k-1}(q^{-1}x) = g_{k-1}(x) \eta_{k-1}(q^{-1}x) \ , \tag{5.55}$$

$$\varphi_k(x) = d_k \frac{B_{k-1}(x)}{B_k(x)} \varphi_{k-1}(q^{-1}x) = \frac{d_k}{g_{k-1}(x)} \varphi_{k-1}(q^{-1}x) \ , \tag{5.56}$$

where the function $g_k(x)$ is given by (2.25). Substituting (5.55, 5.56) into the third relation in (5.52) we obtain finally

$$\eta_{k-1}(x)\varphi_{k-1}^2(x) - \frac{g_{k-1}(qx)}{d_k} \eta_{k-1}(qx) \varphi_{k-1}^2(qx) = \tag{5.57}$$

$$\left(d_k a_{k-1} - a_k + \frac{q^2 d_k B_{k-1}(qx) - g_{k-1}(q^2 x) B_{k-1}(q^2 x)}{(1-q)^2 q^3 x^2} \right) \frac{g_{k-1}(qx)}{d_k^2} \ .$$

Acknowledgement

The authors would like to thank T. Goliński for careful reading of the manuscript and interest in the paper.

References

[1] Chihara, T. S., *An Introduction to Orthogonal Polynomials*, New York: Gordon and Breach, 1978;

[2] Darboux, G., *Sur une proposition relative aux equations lineaires*, C. R. Acad. Sci. Paris, 94 1456, 1882;

[3] Dirac, P.A.M., *Principles of Quantum Mechanics*, Clarendon Press, Oxford, 1947;

[4] Gasper, G. and Rahman, M., *Basic Hypergeometric Series*, Cambridge: Cambridge University Press, 1990;

[5] Goliński, T., Odzijewicz, A., *General difference calculus and its application to functional equations of the second order*, Czechoslovak J. of Phys., **52**, 1219-1224, 2002;

[6] Hahn, W., *Über Orthogonalpolynome die q-Differenzengleichungeg genügen*, Math. Nachr., **2**, 4-34, 1949;

[7] Infeld, L. and Hull, T.E., *The Factorization Method*, Rev. Mod. Phys., **23**, 21-68, 1951;

[8] Koekoek, R. and Swarttouw, R.F., *The Askey–scheme of hypergeometric orthogonal polynomials and its q–analogue*, Report no. 98-17, TUDelft, webpage http://aw.twi.tudelft.nl/koekoek/askey.html, 1998;

[9] de Lange, O.L. and Raab, R.E., Operator Methods in Quantum Mechanics, Claredon Press - Oxford, 1991;

[10] Mielnik, B., Nieto, L.M., Rosas-Ortiz, O., *The finite difference algorithm for higer order supersymmetry*, Phys. Lett. A, **269**, 70-78, 2000;

[11] Miller, W., Jr., *Lie Theory and Special Functions*, Academic Press New York and London, 1968;

[12] Odzijewicz, A., Goliński, T., *Second order functional equations*; webpage http://arXiv:math-ph/0208006, 2002;

[13] Odzijewicz, A., Horowski, H., Tereszkiewicz, A., *Integrable multi-boson systems and orthogonal polynomials*, J. Phys. A: Math. Gen., **34**, 4353-4376, (2001);

[14] Schrödinger, E., Proc. Roy Irish Acad., A **46**, 1940;

[15] Spiridonov, V., *Universal superpositions of coherent states and self–similar potentials*, Phys. Rev. A, **52**, 1909-1935, 1995;

[16] Veselov, A.P., Shabat, A.B., Funct. Anal. Appl., **27**, 1-21, 1993;